

COMPLEX FOLIATIONS AND KÄHLER QCH SURFACES

WŁODZIMIERZ JELONEK

ABSTRACT. The aim of this paper is to describe complex 2-dimensional foliations in a Kähler surface.

1. Introduction. The aim of the present paper is to investigate complex 2-dimensional foliations on connected Kähler surfaces (M, g, J) . Two dimensional complex distribution $\mathcal{D}$ on a Kähler surface defines an opposite almost Hermitian structure I . We prove that if $\mathcal{D}$ is integrable then I is Hermitian if and only if $\mathcal{D}$ is totally geodesic and holomorphic. Also I is Hermitian if $\mathcal{D}$ is conformal and totally geodesic. Hence any complex totally geodesic and conformal foliation is holomorphic. We also prove that the following conditions are equivalent for a complex, 2-dimensional foliation $\mathcal{D}$ on a Kähler surface (M, g, J) : (a) $\mathcal{D}$ is conformal with $L_V g = \theta(V)g$ on $\mathcal{D}^\perp$, (b) $\mathcal{D}$ is quasi holomorphic i.e. $L_\xi J(\mathcal{E}) \subset \mathcal{D}$ for $\xi \in \Gamma(\mathcal{D})$, (c) $2g(\nabla_X \xi, Y) = \theta(\xi)g(X, Y) + J\theta(\xi)\omega(X, Y)$ for $\xi \in \Gamma(\mathcal{D})$ and $X, Y \in \mathcal{E}$ where ω is the Kähler form of (M, g, J) and $d\Omega_I = 2\theta \wedge \Omega_I$ where Ω_I is the Kähler form of (M, g, I) .

A QCH Kähler surface is a Kähler surface admitting a global, 2-dimensional, J -invariant distribution $\mathcal{D}$ having the following property: The holomorphic curvature $K(\pi) = R(X, JX, JX, X)$ of any J -invariant 2-plane $\pi \subset T_x M$, where $X \in \pi$ and $g(X, X) = 1$, depends only on the point x and the number $|X_{\mathcal{D}}| = \sqrt{g(X_{\mathcal{D}}, X_{\mathcal{D}})}$, where $X_{\mathcal{D}}$ is an orthogonal projection of X on $\mathcal{D}$. (see [J-1], [J-2], [J-4] [G-M]). We classify compact, simply connected QCH surfaces for which the opposite almost Hermitian structure I induced by $\mathcal{D}$ is Hermitian and one of the distributions $\ker(J \circ I \pm Id)$ (hence $\mathcal{D}$ or $\mathcal{E} = \mathcal{D}^\perp$) is integrable. We also classify Kähler surfaces (M, g, J) admitting a solution of an equation $H^\phi = g$ for a certain function $\phi \in C^\infty(M)$ where H^ϕ is the Hessian of ϕ .

2. Complex foliations on a Kähler surface. Our notation is as in [K-N]. We start with (see [B-W], [Ch-N]):

Definition 1. A foliation $\mathcal{F}$ on a Riemannian manifold (M, g) is called conformal if

$$L_V g = \alpha(V)g$$

holds on $T\mathcal{F}^\perp$ where α is a one form vanishing on $T\mathcal{F}^\perp$ for any V tangent to the leaves of $\mathcal{F}$. A foliation $\mathcal{F}$ is called homothetic if it is conformal and $d\alpha = 0$.

MS Classification: 53C55, 53C25, 53B35. Key words and phrases: Kähler surface, quasi constant holomorphic sectional curvature, QCH manifold, homothetic foliation, conformal foliation

Definition 2. A complex distribution $\mathcal{D}$ on a complex manifold (M, g, J) is called holomorphic if $L_\xi J(TM) \subset \mathcal{D}$ for any $\xi \in \Gamma(\mathcal{D})$.

In the sequel we shall not make the distinction between a foliation and the distribution of vectors tangent to the leaves of the foliation. A foliation (distribution) $\mathcal{D}$ on a complex manifold (M, J) is called complex if $J(\mathcal{D}) \subset \mathcal{D}$. Complex 2-dimensional foliation (distribution) on a complex surface defines an opposite almost Hermitian structure defined by $I|_{\mathcal{D}} = J|_{\mathcal{D}}, I|_{\mathcal{D}^\perp} = -J|_{\mathcal{D}^\perp}$. If we denote by Ω_J, Ω_I the Kähler forms of (M, g, J) and (M, g, I) respectively then $\Omega_J^2 = -\Omega_I^2$. In the following all the considered distributions $\mathcal{D}, \mathcal{E} = \mathcal{D}^\perp$ are assumed to be complex and 2-dimensional. We shall recall some results from [G-M-1]. Let

$$(2.1) \quad R(X, Y)Z = ([\nabla_X, \nabla_Y] - \nabla_{[X, Y]})Z$$

and let us write

$$R(X, Y, Z, W) = g(R(X, Y)Z, W).$$

If R is the curvature tensor of a QCH Kähler manifold (M, g, J) , then there exist functions $a, b, c \in C^\infty(M)$ such that

$$(2.2) \quad R = a\Pi + b\Phi + c\Psi,$$

where Π is the standard Kähler tensor of constant holomorphic curvature i.e.

$$(2.3) \quad \begin{aligned} \Pi(X, Y, Z, U) &= \frac{1}{4}(g(Y, Z)g(X, U) - g(X, Z)g(Y, U) \\ &+ g(JY, Z)g(JX, U) - g(JX, Z)g(JY, U) - 2g(JX, Y)g(JZ, U)), \end{aligned}$$

the tensor Φ is defined by the following relation

$$(2.4) \quad \begin{aligned} \Phi(X, Y, Z, U) &= \frac{1}{8}(g(Y, Z)h(X, U) - g(X, Z)h(Y, U) \\ &+ g(X, U)h(Y, Z) - g(Y, U)h(X, Z) + g(JY, Z)h(JX, U) \\ &- g(JX, Z)h(JY, U) + g(JX, U)h(JY, Z) - g(JY, U)h(JX, Z) \\ &- 2g(JX, Y)h(JZ, U) - 2g(JZ, U)h(JX, Y)), \end{aligned}$$

and finally

$$(2.5) \quad \Psi(X, Y, Z, U) = -h(JX, Y)h(JZ, U) = -(h_J \otimes h_J)(X, Y, Z, U).$$

where $h_J(X, Y) = h(JX, Y)$.

We say that an almost Hermitian structure I satisfies the second Gray curvature condition if

$$(G2) \quad R(X, Y, Z, W) - R(IX, IY, Z, W) = R(IX, Y, IZ, W) + R(IX, Y, Z, JW).$$

Hence (M, g, I) satisfies the second Gray condition if I preserves the Ricci tensor and W^+ (with respect to the orientation given by I) is degenerate. Note that for a QCH surface (M, g, J) the distribution $\mathcal{D}$ induces an almost Hermitian structure I satisfying the second Gray condition.

3. An almost hermitian structure I induced by complex foliation $\mathcal{D}$. We start with characterizing two dimensional foliations in 4-dimensional Riemannian manifold.

Theorem 1. *Let (M, g) be a Riemann manifold, $\dim M = 4$ and let $\mathcal{D}$ be a two dimensional distribution on M . Let $\{\theta_1, \theta_2, \theta_3, \theta_4\}$ be an orthonormal basis of 1-forms such that $\mathcal{D} = \{X : \theta_3(X) = \theta_4(X) = 0\}$. Let $\omega = \theta_3 \wedge \theta_4$. Then $\mathcal{D}$ is integrable if and only if $d\omega = \phi \wedge \omega$ for a certain $\phi \in \mathcal{A}^1(M)$.*

Proof. $\mathcal{D}$ is integrable if and only if $d\theta_3 = a \wedge \theta_3 + b \wedge \theta_4, d\theta_4 = c \wedge \theta_3 + d \wedge \theta_4$. It follows that if $\mathcal{D}$ is integrable then

$$d\omega = d\theta_3 \wedge \theta_4 - \theta_3 \wedge d\theta_4 = a \wedge \theta_3 \wedge \theta_4 - \theta_3 \wedge d \wedge \theta_4 = (a + d) \wedge \omega.$$

Assume now that $d\omega = \phi \wedge \omega$. Let $d\theta_3 = a \wedge \theta_3 + b \wedge \theta_4 + k\theta_1 \wedge \theta_2, d\theta_4 = c \wedge \theta_3 + d \wedge \theta_4 + l\theta_1 \wedge \theta_2$. Then $d\omega = (a + d) \wedge \omega + k\theta_1 \wedge \theta_2 \wedge \theta_4 - l\theta_3 \wedge \theta_1 \wedge \theta_2$. Hence $(\phi - a - d)\omega = k\theta_1 \wedge \theta_2 \wedge \theta_4 - l\theta_3 \wedge \theta_1 \wedge \theta_2$. Consequently $k = l = 0$ and $\mathcal{D}$ is integrable. $\diamond$

Remark: Let (M, g, J) be a Kähler surface with a Kähler form $\omega = \Omega_J$. Let $\mathcal{D}$ be a complex foliation $J\mathcal{D} \subset \mathcal{D}$. Let us denote by $\mathcal{E} = \mathcal{D}^\perp$ the orthogonal complement of $\mathcal{D}$. Let us define by I an almost Hermitian structure given by $I|_{\mathcal{D}} = J|_{\mathcal{D}}, I|_{\mathcal{E}} = -J|_{\mathcal{E}}$. Then $\Omega_J = \omega_1 + \omega_2, \Omega_I = \omega_1 - \omega_2$ where $\omega_1 \in \bigwedge^2 \mathcal{D}, \omega_2 \in \bigwedge^2 \mathcal{E}$. We have $d\omega_2 = \theta \wedge \omega_2$ where $\theta^\sharp \in \Gamma(\mathcal{D})$. Thus $d\Omega_I = -2\theta \wedge \Omega_I$.

Theorem 2. *Let (M, g, J) be a Kähler surface and $\mathcal{D}$ be a complex foliation, $\dim \mathcal{D} = 2$. Then the almost Hermitian structure I given by $I|_{\mathcal{D}} = J|_{\mathcal{D}}, I|_{\mathcal{D}^\perp} = -J|_{\mathcal{D}^\perp}$ is integrable if and only if $\mathcal{D}$ is holomorphic and totally geodesic.*

Proof. Let us assume that I is integrable. Let $\omega_2 = \theta_3 \wedge \theta_4$ where $\mathcal{D} = \ker \theta_3 \cap \ker \theta_4$. Then $d\omega_2 = \theta \wedge \omega_2$ where $\theta \in \bigwedge^1 \mathcal{D}$. If $\Omega_J = \omega_1 + \omega_2$ then $\Omega_I = \omega_1 - \omega_2$ and $d\Omega_I = -2\theta \wedge \omega_2 = 2\theta \wedge \Omega_I$. Since $\theta^\sharp \in \mathcal{D}$ and I is integrable we have $\nabla_X I = 0$ for $X \in \mathcal{D} = \text{span}\{\theta, J\theta\}$. It follows that $\mathcal{D}$ is totally geodesic. In fact $IX = JX$ for $X, Y \in \mathcal{D}$ implies $I(\nabla_Y X) = J(\nabla_Y X)$. We show that $\mathcal{D}$ is holomorphic. We have to show that $g(\nabla_{JY} X - J\nabla_Y X, Z) = 0$ for arbitrary Y and $Z \in \mathcal{E}$. If Y belongs to $\mathcal{D}$ it is a consequence of the fact that $\mathcal{D}$ is totally geodesic. Let $Y \in \mathcal{E}$. Then $g(\nabla_{JX} \xi, Y) - g(J\nabla_X \xi, Y) = g(\xi, \nabla_{JX} Y + \nabla_X JY)$ for $\xi \in \mathcal{D}, X, Y \in \mathcal{E}$. We have to show that for $X, Y \in \mathcal{E}$ we obtain $\nabla_{JX} Y + \nabla_X JY \in \mathcal{E}$. Taking $Z = JY$ it is equivalent to $-\nabla_{JX} JZ + \nabla_X Z \in \mathcal{E}$ for $X, Z \in \mathcal{E}$. Since $IZ = -JZ$ to $\nabla_X IZ + I(\nabla_X Z) = -J(\nabla_X Z)$. Hence $\nabla_X IZ = -2J(\nabla_X Z|_{\mathcal{D}})$. But I is integrable, hence $\nabla_{IX} IIZ = -2J(\nabla_{IX} IZ|_{\mathcal{D}}) = -2J(\nabla_X Z|_{\mathcal{D}})$. It implies $-\nabla_{JX} JZ + \nabla_X Z \in \mathcal{E}$. Assume now that $\mathcal{D}$ is totally geodesic and holomorphic. Then $\nabla_X I = 0$ for $X \in \mathcal{D}$. If $X, Y \in \mathcal{D}, Z \in \mathcal{E}$ then $\nabla_X Y \in \mathcal{D}, \nabla_X Z \in \mathcal{E}$. It implies $\nabla_X IY = \nabla_X IZ = 0$. Since $\mathcal{D}$ is holomorphic we get $g(\nabla_{JX} \xi - J\nabla_X \xi, Y) = 0$ for $Y \in \mathcal{E}$ and arbitrary X . Let now $Y \in \mathcal{D}, X \in \mathcal{E}$. Then $\nabla_X IY = (J - I)(\nabla_X Y) = 2J(\nabla_X Y|_{\mathcal{E}})$. Note that $\nabla_{IX} IY = -2J(\nabla_{JX} JY|_{\mathcal{E}}) = 2J(\nabla_X Y|_{\mathcal{E}})$. Now assume that $X, Y \in \mathcal{E}$. Then $\nabla_X IY = (-I - J)(\nabla_X Y) = -2J(\nabla_X Y|_{\mathcal{D}})$. Again using that $\mathcal{D}$ is holomorphic we obtain $g(\xi, \nabla_{JX} Y) + g(\xi, \nabla_X JY) = 0$ and consequently $g(\xi, \nabla_X Y - \nabla_{JX} JY) = 0$. Thus $\nabla_X Y|_{\mathcal{D}} = \nabla_{JX} JY|_{\mathcal{D}}$ and $\nabla_{IX} IY = \nabla_X IY$ also in this case. $\diamond$

Theorem 3. *Let (M, g, J) be a Kähler surface. Let us assume that $\mathcal{D}$ is a complex foliation, $\dim \mathcal{D} = 2$ and the structure I defined by $\mathcal{D}$ is integrable. Then*

$$(*) \quad 2g(\nabla_X \xi, Y) = \theta(\xi)g(X, Y) + J\theta(\xi)\omega(X, Y)$$

for every $\xi \in \Gamma(\mathcal{D})$ i $X, Y \in \mathcal{E}$. On the other hand if $\mathcal{D}$ is totally geodesic and $(*)$ is satisfied then I is integrable.

Proof. We have $d\omega_2 = \theta \wedge \omega_2$ and $d\Omega_I = -2\theta \wedge \Omega_I$. Note that $\nabla_X I = 0$ for $X \in \mathcal{D}$. Indeed in the interior of the set $F = \{x : \theta_x = 0\}$ the structure I is Kähler and outside F we have $\text{span}\{\theta^\sharp, J\theta^\sharp\} = \mathcal{D}$. Since I is Hermitian we have $\nabla_\theta I = \nabla_{J\theta} I = 0$. Let $\xi \in \Gamma(\mathcal{D})$. Then $L_\xi - \nabla_\xi = -\nabla \xi$. Note that $\nabla_\xi \omega_1 = \nabla_\xi \omega_2 = 0$. It implies $(L_\xi - \nabla_\xi)\omega_2 = L_\xi \omega_2 = \xi \lrcorner d\omega_2 = \xi \lrcorner (\theta \wedge \omega_2) = \theta(\xi)\omega_2$. Hence $g(J\nabla_X \xi, Y) + g(JX, \nabla_Y \xi) = \theta(\xi)\omega_2(X, Y)$ and $g(\nabla_X \zeta, Y) - g(X, \nabla_Y \zeta) = -\theta(J\zeta)g(JX, Y)$. Since I is Hermitian $\mathcal{D}$ is holomorphic and $g(\nabla_{JX} \xi, Y) + g(JX, \nabla_Y \xi) = \theta(\xi)g(JX, Y)$. Thus

$$g(\nabla_X \xi, Y) + g(X, \nabla_Y \xi) = \theta(\xi)g(X, Y) \text{ for any } X, Y \in \mathcal{E}. \text{ Consequently}$$

$$2g(\nabla_X \xi, Y) = \theta(\xi)g(X, Y) + J\theta(\xi)\omega(X, Y).$$

On the other hand if $\mathcal{D}$ is totally geodesic and $(*)$ is satisfied then $\mathcal{D}$ is holomorphic and I is integrable. $\diamond$

Theorem 4. *Let $\mathcal{D}$ be a 2-dimensional, complex foliation on a Kähler surface (M, g, J) . Then $\mathcal{E} = \mathcal{D}^\perp$ is integrable if and only if I is almost Kähler.*

Proof. If $\mathcal{E}, \mathcal{D}$ are integrable then $d\omega_1 = \phi \wedge \omega_1, d\omega_2 = \psi \wedge \omega_2$ and $d\omega_1 = -d\omega_2$ hence $d\omega_1 = d\omega_2 = 0$. On the other hand if $d\omega_1 = d\omega_2 = 0$ then $\mathcal{D}, \mathcal{E}$ are integrable. $\diamond$

Theorem 5. *Let (M, g, J) be a Kähler surface and $\mathcal{D}$ be a complex distribution $J\mathcal{D} \subset \mathcal{D}$ and I be an almost Hermitian structure defined by $\mathcal{D}$. Then $\mathcal{D} \subset \ker \nabla I = \{X : \nabla_X I = 0\}$ if and only if $\mathcal{D}$ is totally geodesic.*

Proof $\Rightarrow$ Let $X \in \mathcal{D}$ then $IX = JX$ and for $Y \in \mathcal{D}$ we get $I\nabla_Y X = J\nabla_Y X$ and consequently $\nabla_Y X \in \mathcal{D}$.

$\Leftarrow$ Let $\mathcal{D}$ be totally geodesic. Then $\nabla_X Y \in \mathcal{D}(\mathcal{E})$ if $Y \in \mathcal{D}(\mathcal{E})$ and $X \in \mathcal{D}$. It implies $\nabla_X IY + I\nabla_X Y = \pm J\nabla_X Y$ and $\nabla_X IY = 0$ for $Y \in \mathcal{D}(Y \in \mathcal{E})$. $\diamond$

Remark. Let (M, g, J) be a 3-symmetric Kähler surface (see [K]). Then (M, g, J) is a QCH surface and it admits strictly almost Kähler opposite structure I . If we denote by $\mathcal{D}$ the Kähler nullity of I then (M, g, J) is a QCH with respect to $\mathcal{D}$ and $\mathcal{D}$ is totally geodesic. Hence there exist totally geodesic complex foliations on Kähler surfaces for which the almost Hermitian structure I induced by $\mathcal{D}$ is not integrable.

Lemma. *Let us assume that $J\mathcal{D} = \mathcal{D}$ and $\mathcal{E} = \mathcal{D}^\perp$. If $\omega_1 \in \bigwedge^2 \mathcal{D}, \omega_2 \in \bigwedge^2 \mathcal{E}$ and $\omega = \omega_1 + \omega_2$ where ω is a Kähler form of (M, g, J) then $\nabla_X \omega_2(Y, Z) = 0$ for $X \in \mathcal{D}, Y, Z \in \mathcal{E}$.*

Proof We have $\omega_2(Y, Z) = g(JY, Z)$. Note also that $\omega_2(Y, Z) = g(JY, Z)$ if $Z \in \mathcal{E}$ and Y is arbitrary. Thus

$$\begin{aligned} \nabla_X \omega_2(Y, Z) + \omega_2(\nabla_X Y, Z) + \omega_2(Y, \nabla_X Z) &= g(J\nabla_X Y, Z) + g(JY, \nabla_X Z) \\ &= \omega_2(\nabla_X Y, Z) + \omega_2(Y, \nabla_X Z) \end{aligned}$$

◇

Corollary 1. *If $\mathcal{D}$ is complex and integrable then*

$$g(\nabla_X \xi, Y) - g(\nabla_Y \xi, X) = J\theta(\xi)g(JX, Y)$$

for $\xi \in \Gamma(\mathcal{D})$ and $X, Y \in \mathcal{E}$ where $d\Omega_I = -2\theta \wedge \Omega_I$. For $X, Y \in \Gamma(\mathcal{E})$ we have $[X, Y]_{|\mathcal{D}} = -J\theta^\# \omega(X, Y)$.

Proof. We have $\Omega_J = \omega_1 + \omega_2$, $\Omega_I = \omega_1 - \omega_2$ and $d\omega_2 = \theta \wedge \omega_2$ where $\theta^\# \in \Gamma(\mathcal{D})$. Hence $(L_\xi - \nabla_\xi)\omega_2 = L_\xi\omega_2 - \nabla_\xi\omega_2 = \theta(\xi)\omega_2 - \nabla_\xi\omega_2$. It follows that for $Y, Z \in \mathcal{E}$ we get $\omega_2(\nabla_X \xi, Y) + \omega_2(X, \nabla_Y \xi) = \theta(\xi)\omega_2(X, Y)$ and $g(\nabla_X J\xi, Y) - g(X, \nabla_Y J\xi) = \theta(\xi)g(JX, Y)$. Thus $g(\nabla_X \xi, Y) - g(X, \nabla_Y \xi) = J\theta(\xi)g(JX, Y)$.

Hence $(**)[X, Y]_{|\mathcal{D}} = -J\theta^\# \omega(X, Y)$. ◇

Remark Let us note that $d\theta(X, Y) = 0$ for $X, Y \in \mathcal{D}$ and $d\theta(X, Y) = 0$ for $X, Y \in \mathcal{E}$. In fact from $(**)$ it follows for $X, Y \in \mathcal{E}$. Next $d\theta \wedge (\theta_1 \wedge \theta_2 - \theta_3 \wedge \theta_4) = 0$. Hence the result follows.

Theorem 6. *The following conditions are equivalent for a complex, 2-dimensional foliation $\mathcal{D}$ on a Kähler surface (M, g, J) :*

- (a) $\mathcal{D}$ is conformal
- (b) $\mathcal{D}$ is quasi holomorphic i.e. $L_\xi J(\mathcal{E}) \subset \mathcal{D}$ for $\xi \in \Gamma(\mathcal{D})$
- (c) $2g(\nabla_X \xi, Y) = \theta(\xi)g(X, Y) + J\theta(\xi)\omega(X, Y)$ for $\xi \in \Gamma(\mathcal{D})$ and $X, Y \in \mathcal{E}$ where $d\Omega_I = -2\theta \wedge \Omega_I$.

Proof. (a) $\Rightarrow$ (b), (c) Note that $g(\nabla_X \xi, Y) - g(\nabla_Y \xi, X) = J\theta(\xi)g(JX, Y)$ and $g(\nabla_X \xi, Y) + g(\nabla_Y \xi, X) = \phi(\xi)g(X, Y)$. Summing up

$$2g(\nabla_X \xi, Y) = J\theta(\xi)g(JX, Y) + \phi(\xi)g(X, Y).$$

Thus $g(\nabla_{JX} \xi, JY) = g(\nabla_X \xi, Y)$. It is also easily seen that $\phi = \theta$.

(b) $\Rightarrow$ (a) If $\mathcal{D}$ is quasi holomorphic then from

$$g(\nabla_X \xi, JY) - g(\nabla_{JY} \xi, X) = J\theta(\xi)g(JX, JY)$$

we get

$$-g(\nabla_X J\xi, Y) - g(\nabla_Y J\xi, X) = -\theta(J\xi)g(X, Y)$$

and consequently

$$g(\nabla_X \xi, Y) + g(\nabla_Y \xi, X) = \theta(\xi)g(X, Y)$$

and $\mathcal{D}$ is conformal.

Corollary 2. *If $\mathcal{D}$ is a complex foliation on a Kähler surface (M, g, J) which is conformal and totally geodesic then the opposite almost hermitian structure is Hermitian. Hence conformal and totally geodesic foliation $\mathcal{D}$ is holomorphic.*

Proof. Since $\mathcal{D}$ is conformal is quasi holomorphic. Since it is totally geodesic it is holomorphic. Hence the result follows from Th.2. ◇

Theorem 7. *Let (M, g, J) be a compact simply connected QCH Kähler surface. If I is Hermitian and one of the distributions $\ker(I \circ J \pm Id)$ is integrable and then (M, g, J) is globally of Calabi type or a product of Riemann surfaces.*

Proof. Let $\mathcal{D} = \ker(J \circ I - Id)$ be integrable. Then $\mathcal{D}$ is totally geodesic homothetic foliation. In fact since I is integrable we have $d\theta^- = 0$ and since M is compact $d\theta = 0$ (see [A-G]). Note that that homothetic foliations are classified locally in [Ch-N]. Thus if we do not assume that M is simply connected but is compact it follows that locally is isometric to a Calabi type surface or a product of Riemann surfaces. In the compact simply connected case the result follows from [J-3]. $\diamond$

Theorem 8. *Let $\mathcal{D}$ be a complex totally geodesic and holomorphic foliation on a Kähler surface (M, g, J) . Then $\mathcal{D}$ is conformal, $2g(\nabla_X \xi, Y) = |\theta|^2 g(X, Y)$ where $\theta = g(\xi, \cdot)$ and $d\bar{\omega} = -2\theta \wedge \bar{\omega}$, $L_\eta g = \theta(\eta)g$ on $\mathcal{D}^\perp$ for $\eta \in \Gamma(\mathcal{D})$. If $d\theta^- = 0$ then (M, g, J) is a QCH Kähler surface whose negative almost Hermitian G_2 structure $\bar{J}$ is Hermitian. On the other hand if (M, g, J) is a QCH Kähler surface with integrable distribution $\mathcal{D}$ and Hermitian structure $\bar{J}$ then $\mathcal{D}$ is a totally geodesic conformal holomorphic foliation with $d\theta^- = 0$. $\diamond$*

Proof We have for $X, Y \in \Gamma(\mathcal{D}^\perp)$ $2g(\nabla_X \xi, Y) = |\theta|^2 g(X, Y)$ thus for $Z \in \Gamma(\mathcal{D}^\perp)$ we obtain

$$2g(\nabla_Z \nabla_X \xi, Y) + 2g(\nabla_X \xi, \nabla_Z Y) = \\ Z|\theta|^2 g(X, Y) + |\theta|^2 g(\nabla_Z X, Y) + |\theta|^2 g(X, \nabla_Z Y)$$

If $d\theta^- = 0$ then $d\theta(JX, JY) = -d\theta(X, Y)$. Hence

$$d\theta(\xi, X) = -\frac{1}{2}X|\theta|^2, d\theta(J\xi, X) = -\frac{1}{2}JX|\theta|^2$$

and $\nabla_Z Y = -\frac{1}{2}g(Z, Y)\xi - \frac{1}{2}\omega(Z, Y)J\xi + \nabla_Z Y_{\mathcal{D}^\perp}$. Consequently

$$2g(\nabla_Z \nabla_X \xi, Y) = \frac{1}{2}g(Z, Y)X|\theta|^2 + \frac{1}{2}\omega(Z, Y)JX|\theta|^2 \\ + Z|\theta|^2 g(X, Y) + |\theta|^2 g(\nabla_Z X, Y)$$

Hence

$$4g(R(Z, X)\xi, Y) = -X|\theta|^2 g(Z, Y) + Z|\theta|^2 g(X, Y) + \\ \omega(Z, Y)JX|\theta|^2 - \omega(X, Y)JZ|\theta|^2 = 0.$$

It follows that $(M, g, \bar{J})$ is a G_3 Gray manifold, and since $\bar{J}$ is Hermitian it follows that $(M, g, \bar{J})$ is a G_2 manifold and (M, g, J) is a QCH manifold. We have assumed here that $\theta \neq 0$ but if $\theta = 0$ then (M, g, J) is a product hence also a QCH manifold. $\diamond$

Now we consider Kähler QCH surfaces which are semi-symmetric (i.e. $R.R = 0$) with Hermitian structure $\bar{J}$ and such that $(M, g, \bar{J})$ is of Gray type (G_1). These are semi-symmetric surfaces foliated by two dimensional Euclidean space (see [J-5]). Let $\omega \in \bigwedge^2 \mathcal{D}$ be as above. We have $\nabla_\xi \omega = 0$ and $-2\nabla_X \omega = -JX \wedge \theta + X \wedge J\theta$ for $X \in \mathcal{D}^\perp$. Hence $\nabla_X \nabla_\xi \omega = 0$,

$$-2\nabla_\xi \nabla_X \omega = -J(\nabla_\xi X) \wedge \theta - JX \wedge \nabla_\xi \theta + \nabla_\xi X \wedge J\theta + X \wedge J\nabla_\xi \theta.$$

Note that $\nabla_X \xi = \frac{1}{2}|\theta|^2 X + \frac{1}{2}X \ln |\theta|^2 \xi + \frac{1}{2}JX \ln |\theta|^2 J\xi$, and

$$\begin{aligned} -2\nabla_{[X, \xi]}\omega &= -J[X, \xi]_\perp \wedge \theta + [X, \xi]_\perp \wedge J\theta = \\ &= -J\nabla_X \xi_\perp \wedge \theta + J\nabla_\xi X \wedge \theta + \frac{1}{2}|\theta|^2 X \wedge J\theta - \nabla_\xi X \wedge J\theta = \\ &= -\frac{1}{2}|\theta|^2 JX \wedge \theta + J\nabla_\xi X \wedge \theta + \frac{1}{2}|\theta|^2 X \wedge J\theta - \nabla_\xi X \wedge J\theta \end{aligned}$$

Hence

$$-2R(\xi, X) \cdot \omega = JX \wedge \nabla_\xi \theta - X \wedge J\nabla_\xi \theta + \frac{1}{2}|\theta|^2 JX \wedge \theta - \frac{1}{2}|\theta|^2 X \wedge J\theta$$

Consequently (note that $R(\xi, X) = 0$) $\nabla_\xi \theta + \frac{1}{2}|\theta|^2 \xi = 0$ i $\nabla_\xi \xi = -\frac{1}{2}|\theta|^2 \xi$ and $\nabla_{J\xi} \xi = \frac{1}{2}|\theta|^2 J\xi$. We also have $\xi|\theta|^2 = -|\theta|^4$ and $[\xi, J\xi] = -|\theta|^2 J\xi$

Now we shall show that $\eta = \frac{1}{|\theta|^2} \xi$ is a holomorphic vector field and $J\eta$ is a holomorphic divergence free vector field.

Note that $\nabla_X \eta = \frac{1}{2}X - \frac{1}{2}X \ln |\theta|^2 \eta + \frac{1}{2}JX \ln |\theta|^2 J\eta$ for $X \in \mathcal{D}^\perp$. Hence $\nabla_{JX} \eta = J\nabla_X \eta$.

Similarly $\nabla_{J\xi} \eta = \frac{1}{2}J\xi$ and $\nabla_\xi \eta = \frac{1}{2}\xi$. Hence $\nabla_X \eta = \frac{1}{2}X$ for $X \in \mathcal{D}$. It is also clear that $\nabla_{JX} \eta = J\nabla_X \eta$ if $X \in \mathcal{D}$ and $\eta \in \mathfrak{hol}(M)$. We also have $\nabla_X J\eta = \frac{1}{2}JX - \frac{1}{2}X \ln |\theta|^2 J\eta - \frac{1}{2}JX \ln |\theta|^2 \eta$ for $X \in \mathcal{D}^\perp$ and $\nabla_X J\eta = \frac{1}{2}JX$ for $X \in \mathcal{D}$. Consequently $\text{div} J\eta = 0$.

If $d\theta = 0$ then $\theta = -d \ln |\theta|^2$ and consequently on semi-symmetric surfaces foliated by two dimensional Euclidean space of Calabi type we have $\theta \neq 0$ on the whole of M .

Theorem 9. *Let (M, g, J) be a Kähler surface. Then the following conditions are equivalent:*

- (1) *There exists a vector field ξ such that $\nabla \xi = cI, c \in \mathbb{R} - \{0\}$*
- (2) *There exists a vector field η such that $\nabla \eta = cJ, c \in \mathbb{R} - \{0\}$*
- (3) *There exists a function $\phi \in C^\infty(M)$ such that $H^\phi = c g$ where H^ϕ is the Hessian of ϕ and $c \in \mathbb{R} - \{0\}$,*
- (4) *(U, g, J) is a semi-symmetric Kähler surface of Calabi type where U is an open dense subset of M ,*
- (5) *There exists an open and dense subset U of M such that (U, g, J) is locally isometric to*

$$g = z g_\Sigma + \frac{1}{Cz} dz^2 + Cz(dt + \alpha)^2$$

where (Σ, g_Σ) is a Riemannian surface with area form ω_Σ and $d\alpha = \omega_\Sigma$. The Kähler form of (U, g, J) is $\Omega = z\omega_\Sigma + dz \wedge (dt + \alpha)$.

Proof. Let us assume that (1) holds. It is easy to see that (1),(2) are equivalent and (3) implies (1).

Note that $\nabla_X J\xi = cJX$ so we can take $\eta = J\xi$. Let us define the distribution $\mathcal{D} = \text{span}\{\xi, J\xi\}$. We can assume that $c = \frac{1}{2}$. It is clear that ξ is a holomorphic vector field and $J\xi$ is a holomorphic Killing vector field. Note that $\xi, J\xi$ are different from zero on an open dense subset U of M . What is more if $T = \nabla J\xi = \frac{1}{2}J$ then $R(X, \xi)Y = \nabla T(X, Y) = 0$. Hence $R(X, Y)\xi = R(X, Y)J\xi = 0$ and

$$R(X, JX, JX, X) = \|X_{\mathcal{D}^\perp}\|^4 K(\mathcal{D}^\perp)$$

where $K(\mathcal{D}^\perp)$ is a sectional curvature of the distribution $\mathcal{D}^\perp$. It follows that (M, g, J) is a QCH Kähler surface i $R = c\Psi$ and $R.R = 0$. The distribution $\mathcal{D}$ is totally geodesic in particular is integrable. Since $L_{J\xi}g = 0, L_\xi g = g$ it follows that $\mathcal{D}$ is a complex conformal foliation and the almost Hermitian structure I determined by $\mathcal{D}$ is Hermitian. Note that $L_\eta g = \theta(\eta)g$ on $\mathcal{D}^\perp$ for $\eta \in \Gamma(\mathcal{D})$ where $\theta^\# = \frac{1}{|\xi|^2}\xi$. Hence $|\theta| = \frac{1}{|\xi|}$. Vector field $\xi = \frac{1}{|\theta|^2}\theta^\#$ is holomorphic. One can easily verify that $d\theta = 0$ since $X|\theta|^2 = 0$ for $X \in \mathcal{D}^\perp$ since $\nabla_X \xi = \frac{1}{2}X - \frac{1}{2}X \ln |\theta|^2 \xi + \frac{1}{2}JX \ln |\theta|^2 J\xi$ for $X \in \mathcal{D}^\perp$. Hence (M, g, J) is a semi-symmetric Kähler surface of Calabi type. Note that since $\theta = -d \ln |\theta|^2$ on U it follows that if the distribution $\mathcal{D}$ extends over the whole of M and consequently θ is defined on M then $\theta \neq 0$ on the whole of M and consequently $U = M$. Note that the function $\phi = \frac{1}{|\theta|^2}$ satisfies (3) and the field $\xi = \nabla \phi$ satisfies (1).

On the other hand if (M, g, J) is a semi-symmetric Kähler surface of Calabi type then vector field $\xi = \frac{1}{|\theta|^2}\theta^\#$ satisfies $\nabla_X \xi = \frac{1}{2}X$. Semi-symmetric Kähler surfaces of Calabi type are classified in [J-5] which gives equivalence (4) $\Leftrightarrow$ (5). $\diamond$

Remark. It is known (see [T])that the only complete Kähler surface satisfying (3) is a Euclidean space $(\mathbb{C}^2, can)$ with standard metric *can*. Let $\{e_1, e_2, e_3, e_4\}$ be a standard orthonormal basis of $\mathbb{C}^2$, $Je_1 = e_2, Je_3 = e_4$. Then $\xi = x_1e_1 + y_1e_2 + x_2e_3 + y_2e_4, J\xi = x_1e_2 - y_1e_1 + x_2e_4 - y_2e_3$ where $z_1 = x_1 + iy_1, z_2 = x_2 + iy_2$ are standard complex coordinates on $\mathbb{C}^2$ and $\phi = \frac{1}{2}(x_1^2 + y_1^2 + x_2^2 + y_2^2)$. In this case we have $U = \mathbb{C}^2 - \{0\}$ and the totally geodesic complex foliation $\mathcal{D} = span\{\xi, J\xi\}$ defines on U a Hermitian conformally Kähler structure I which does not extend to the whole of $\mathbb{C}^2$ and $\xi(0) = 0$. Hence $(\mathbb{C}^2 - \{0\}, can)$ is a semi-symmetric surface of Calabi type (clearly $R = 0$ in this case). Note that orthogonal complex structures on domains of $\mathbb{C}^2$ are described in [S-V].

References.

- [A-G] V. Apostolov, P. Gauduchon The Riemannian Goldberg-Sachs Theorem Internat. J. Math. vol.8, No.4, (1997),421-439
- [B-W] P. Baird and J. Wood *Harmonic morphisms between Riemannian manifolds* London Mathematical Society Monographs 29 (Oxford University Press, Oxford 2003)
- [Bes] A. L. Besse *Einstein manifolds*, Ergebnisse, ser.3, vol. 10, Springer-Verlag, Berlin-Heidelberg-New York, 1987.
- [Ch-N] S.G.Chiossi and P-A. Nagy *Complex homothetic foliations on Kähler manifolds* Bull. London Math. Soc. 44 (2012) 113-124.
- [G-M] G.Ganchev, V. Mihova *Kähler manifolds of quasi-constant holomorphic sectional curvatures*, Cent. Eur. J. Math. 6(1),(2008), 43-75.
- [J-1] W. Jelonek, Compact holomorphically pseudosymmetric Kähler manifolds Coll. Math.117,(2009),No.2,243-249.
- [J-2] W.Jelonek *Kähler manifolds with quasi-constant holomorphic curvature*, Ann. Glob. Anal. and Geom, vol.36, p. 143-159,(2009)
- [J-3] W. Jelonek, *Kähler manifolds with homothetic foliation by curves* arxiv

[J-4] W. Jelonek *Kähler surfaces with quasi-constant holomorphic curvature* to appear in Glasgow Math.J.

[J-5] W. Jelonek *Semi-symmetric Kähler surfaces* arxiv

[K-N] S. Kobayashi and K. Nomizu *Foundations of Differential Geometry*, vol.2, Interscience, New York 1963

[K] O. Kowalski *Generalized symmetric spaces* Lecture Notes in Math. 805, Springer, New York, 1980.

[S-V] Salamon S., Viaclovsky J., *Orthogonal complex structures on domains in $\mathbb{R}^4$* Math. Ann. 343, 853-899 (2009)

[T] Y. Tashiro, Complete Riemannian manifolds and some vector fields, Trans. Amer. Math. Soc. 117(1965) 251 275

Institute of Mathematics
Cracow University of Technology
Warszawska 24
31-155 Kraków, POLAND.
E-mail address: wjelon@pk.edu.pl